arXiv:1412.5859v1 [physics.ao-ph] 18 Dec 2014

Investigation of carbon dioxide phase shift possibility under extreme Antarctic winter conditions

V.M.Vashchenko, Ye.A. Loza

State ecological academy
of post graduate education and management
Ukraine, Kyiv, V.Lypkivskogo str., 35
Loza@bmyr.kiev.ua

Abstract

The Antarctic winter atmosphere minimal temperature and pressure series reveal that CO_2 phase shift (deposition) is possible in some extreme cases, even leading to possible CO_2 snow phenomenon at Vostok Antarctic station and in other near South Pole regions. A hypothesis has been formulated that stable CO_2 snow cover might have formed in Earth past which may influence interpretation of glacial chronology records. This effect may also manifest in other minor gases. Its global climate role is discussed.

Keywords: *carbon dioxide, phase shift, Antarctic, atmosphere, CO_2 snowflake.*

1 Introduction

There are two main gases in the Earth atmosphere - oxygen and nitrogen. However, minor gases, like H_2O and CO_2 play a very important role in Earth global climate [1]. The mole fractions and phase shifts temperatures of gases with mole fraction over 0.01% are given in Table 1.

Oxygen and nitrogen have boiling and melting (freezing) points at temperatures of $-180\ ^oC$ and lower under normal atmospheric pressure. And so does argon.

However H_2O experiences phase shifts in the atmosphere with well known consequences as rain and snow (starting from liquid or solid aerosol formation), having key influence on Earth global climate. Moreover the other

Table 1: Main atmospheric gases with mole fraction over 0.01% [3] and phase change temperatures at normal conditions.

Gas	**Melting (sublimation)**		**Boiling**		**Mole fraction**
N_2	-210 ^oC	63 ^oK	-196 ^oC	77 ^oK	78.08%
O_2	-183 ^oC	90 ^oK	-219 ^oC	54 ^oK	20.95%
H_2O	0 ^oC	273 ^oK	+100 ^oC	373 ^oK	≈0.5% average
Ar	-189 ^oC	84 ^oK	-186 ^oC	87 ^oK	0.93%
CO_2	-79 ^oC	194 ^oK	n/a	n/a	0.04%
other	n/a	n/a	n/a	n/a	<0.001% each

aerosol particles may experience phase shifts under atmosphere temperature change [2]. However, this phenomenon has not yet been well studied.

This phenomenon may significantly influence the atmosphere state and the results of atmosphere investigations by optical methods and must be included in climate models.

Carbon dioxide cannot experience boiling/precipitation under any extreme atmospheric pressure values in natural conditions, requiring around 5200 mb with ever-observed natural maximum of 1100 mb.

In this paper we investigate CO_2 behavior at extreme cold temperatures. The registered lowest temperature record for the last century is $-89.2\ ^oC$ ("Vostok" station, Antarctic, 1983). As far as CO_2 sublimation/deposition temperature at 1000 mb pressure is $-78.5\ ^oC$ it is plausible that under such extreme conditions we may indeed witness CO_2 snow phenomenon at Earth South Pole.

2 Atmosphere temperature ranges and other conditions

First of all we should mention lowest temperature records close to CO_2 sublimation/deposition temperature found on Earth and summarized in table 2.

We should also note that at different heights the atmosphere temperature is usually lower than that on the surface (e.g. by 45 to 75 degrees for height of ozone maximum which is 30-40 km). The pressure also reduces (e.g. from 1000 to 20-100 mb for the same height of ozone maximum). However, the real stratosphere zonal minimal temperatures at 70 mb heights measured by remote sounding is around $-91\ ^oC$, sometimes reaching $-93\ ^oC$ in the Antarctic [7]. We must also stress that all the atmospheric gases usu-

Table 2: Notable world low temperature records.

Location	**Lowest temperature record**	**Date of the record**	**Site height**	**Pressure (approximately)**
South Pole	-82.8 ^oC	23 Jun 1982	2800 m	800 mb
Vostok station [4]	-89.2 ^oC -91 ^oC unconfirmed	21 Jul 1983 winter 1997	3488 m	620 mb [5]
Oimekon (Siberia)[6]	-67.7 ^oC -76 ^oC disputed	15 Jan 1885	800 m	900 mb

ally experience "supercooled" state. The temperatures required to form the snowflake may be by up to 40 degrees lower than the melting point.

The temperature anomalies may appear at some local points leading to appearance of some zones where CO_2 snow may form and then transport to areas with warmer conditions. The aerosol particles formed in the upper atmosphere under lower temperatures may quickly evaporate at the lower heights. I.e. we may be able observe quasi-stable snow forming and falling at a limited heights range.

Another important condition for aerosol and snow formation is availability of nucleation centers.

Concentrations of small, intermediate and large positive ions in Antarctica was estimated to be 2 to 6 $\times$ 10^2 cm^{-3}, 7 to 30 $\times$ 10^2 cm^{-3}, and from 5 to 12 $\times$ 10^3 cm^{-3} respectively [8] at near-shore station Maitri which corresponds to average over-sea concentrations.

OH concentration in the Antarctic is higher than that at equator and constitutes about 2 $\times$ 10^6 molecules per cubic centimeter of air [9]. Another favorable factor is that ionization radiation level at South Pole is higher due to Earth magnetic field properties.

Well-known H_2O aerosols and snowflakes may also be nucleation centers for CO_2 deposition, and therefore increased quantity of CO_2 in regular snow is another possible manifestation of CO_2 phase shift under extremely cold temperatures. I.e. CO_2 aerosol forms and links to H_2O snowflake, increasing overall CO_2 concentration in the snow layer and leading to ambiguity error in CO_2 glacial chronology records interpretation.



Figure 1: CO_2 phase diagram, based on [10].

Table 3: CO_2 sublimation point pressure dependency based on fig.1

Pressure	**Sublimation point temperature**
1000 mb (sea-level)	$-78.5\ ^oC$
800 mb (South Pole)	$-78.8\ ^oC$
600 mb (Vostok station)	$-82.4\ ^oC$
70 mb (Stratosphere)	$-104.8\ ^oC$

3 CO_2 phase shift

As far as this article is accented at Antarctic atmosphere, we should also mention more usual temperatures for "Vostok" station region. These are $-68.0\ ^oC$ average winter temperature and $-31.9\ ^oC$ average summer temperature. The average minimum temperature is $-71.6\ ^oC$ and average maximum temperature is $-50.3\ ^oC$ (with record at $-12.2\ ^oC$).

As stated above, the stratosphere temperature at 70 mb height does not reach temperatures lower than $-93\ ^oC$ [7]. Therefore, according to table 3 snow formation is impossible at these heights even at the record cold temperatures.

However, sublimation point temperature $-82.4\ ^oC$ at Vostok station, while exceeding average minimum temperature of $-71.6\ ^oC$, is quiet enough for extremal minimum temperatures, such as the 1983 temperature record

of $-89.2\ ^oC$ with difference about 7 degrees.

Let's also note that South Pole with record minimal temperature of $-82.8\ ^oC$ and corresponding sublimation point temperature $-78.8\ ^oC$ is also a candidate for deposition of CO_2 with difference about 4 degrees.

Therefore, we may expect CO_2 phase shift as a rare event during extremely cold Antarctic winters.

4 CO_2 snowflakes

Another problem to consider is the detection possibility and characteristics of the CO_2 snowflakes. First of all, we should note that a regular CO_2 snowflakes size is estimated between 4 and 22 microns [11].

Moreover, all the theoretical and experimental research considering CO_2 snowflakes was made for Martian-like conditions (temperatures down to $-140\ ^oC$). By analogy with H_2O snowflakes we may expect even smaller sizes of CO_2 snowflakes formed under conditions near to sublimation temperature. Therefore, the snowflake size should not exceed 5 microns. I.e. we should not witness regular snowflakes, but rather unseen by eye and even simple microscopic devices several microns size aerosol.

The globally averaged annual H_2O precipitation is 990 millimeters [13]. According to global average CO_2 concentration 0.04% comparing to global average water vapor concentration around 1% we may expect that CO_2 snow quantity generated during extreme cold period (let's consider it 24 hours) would hardly exceed 0.1 mm.

However, considering CO_2 snowflake size of 5 microns we may expect about 20 layers of snowflakes in the snow cover. In this case we should expect a tiny carbon dioxide rich layer of usual H_2O snow.

Therefore CO_2 snowflakes may be definitely detected by: 1) electronic microscopy, leading to difficulty of "catching" and "transporting" the snowflakes; 2) by remote optical method such as spectropolarimetry [12] leading to ambiguous results and 3) by nuclear-activation method on aerosol filters creating "transporting" and "storage" difficulty.

On the other hand super-cold chemical detection methods may be used e.g. detecting CO_2 gas level in sublimation camera after 10-30 microns filter, thou the average CO_2 snow quantity will be low, requiring high method sensitivity. Due to CO_2 gas density higher than that of average atmosphere density, the device filter should be placed at small angle (no greater than 45^o) with CO_2 accumulation & sublimation camera located below the filter. Additional thermocouple is required to control the filter temperature not ex-

ceeding CO_2 sublimation temperature and sublimation camera temperature slightly exceeding this limit.

One more point to consider is that CO_2 snow is very likely to be mixed with regular H_2O snow both in the air and in the snow cover. The pores in the upper snow layer about 0.1 mm wide will accumulate CO_2 snow quickly sublimating it due to higher internal temperature of the H_2O snow layer.

On the other hand, in air CO_2 snow and H_2O snow in the air are easily separated by simple aerosol filters because the majority of CO_2 snowflakes will be much smaller than most of the H_2O snowflakes.

5 Possibility of carbon dioxide stable snow cover at South Pole

In case Earth global temperature (or just regional Antarctic temperature) drops by about 10 degrees, the CO_2 snow in Antarctic winter would appear on regular basis, followed by stable CO_2 winter snow in case of temperature reduction by 14 degrees.

As seen by EPICA and other glacial chronology records the Antarctica region show that Earth climate was close to these limits during some cold periods (see fig. 2).

Moreover temperature never actually reduces lower than 10 degrees than present average, as if there is a stable limit. This limit may be CO_2 stable snow cover formation. This effect may caused by neglecting stable CO_2 snow cover in ice records leading to significant temperature overestimate during such periods.

Different concentration of oxygen isotopes may be caused by admixture of CO_2 snow to regular H_2O snow therefore causing misinterpretation of temperature in cryochronological records. And steady minimal temperature increase in the past may be caused by paleo CO_2 diffusing and leaving its original deposition layer and even diffusing to the surface and leaving snow cover with time.

This causes some degree of enrichment of regular snow with O and C isotopic compound different from normal H_2O snow at the given temperature [15, 16].

6 Minor gases

In addition to CO_2 the following minor gases (with mole fraction over 10^{-7} %) may experience phase shifts in the atmosphere during in-year tem-



Figure 2: Vostok Ice Core temperature records approximately adapted to current minimal winter temperatures based on [14].

perature variations (Table 4 does not present a complete list).

While such gases as O_3 and Xe may hardly form phase shifts even under extreme temperature records, thou some tiny aerosol particles may form locally, N_2O may experience a phase shift in the atmosphere at extreme conditions. Moreover, HF, Hg and NH_3 have phase shift temperatures that stably appear in a large fraction of the Earth surface. And NO_2, like H_2O, may experience two phases shifts in a large fraction of the Earth surface.

Thou their concentration is far smaller than that of CO_2, such phase shifts definitely influence global climate by stabilizing respective temperatu-

Table 4: Minor gases molar concentrations, melting and boiling temperatures.

Gas	**Melting (sublimation)**		**Boiling**		**Mole fraction**
O_3	-192 ^oC	81 ^oK	-112 ^oC	161 ^oK	10^{-5} %
Xe	-112 ^oC	161 ^oK	-108 ^oC	165 ^oK	10^{-5} %
N_2O	-91 ^oC	182 ^oK	-88 ^oC	185 ^oK	10^{-4} %
NO_2	**-11** ^oC	**262** ^oK	**21** ^oC	**294** ^oK	10^{-5} %
HF	-84 ^oC	189 ^oK	**20** ^oC	**293** ^oK	less than 10^{-5} %
Hg	**-30** ^oC	**243** ^oK	357 ^oC	630 ^oK	less than 10^{-5} %
NH_3	-78 ^oC	195 ^oK	**-33** ^oC	**240** ^oK	less than 10^{-5} %

res at certain phase shifts temperature values and by changing atmosphere chemistry and radiative parameters. Moreover, formation of such quasi-seasonal atmospheric aerosol changes must be included in atmospheric models used for Earth atmosphere remote measurements.

However, investigation of these phase shift manifestation is outside of the scope of this article.

7 Conclusions

1. An overview of possibility of phase shifts of minor gases in the Earth atmosphere was made with accent at carbon dioxide. It was found that CO_2 may form solid aerosol and even microscopic snowflakes during extreme Antarctic winter conditions at South Pole and Vostok station.

2. A hypothesis has been formulated that stable CO_2 snow cover might have formed in Earth past which may influence interpretation of glacial chronology records.

3. A brief analysis of other minor gases phase shifts conditions has been made. It was found that N_2O, HF, Hg, NH_3 and NO_2 may also experience phase shifts in Earth atmosphere temperature range. Seasonal changes in atmosphere aerosol composition and aggregate state may impact the results of remote optical observations of the Antarctic atmosphere and should be included in climate models.

References

[1] Русов В.Д., Глушков А.В., Ващенко В.Н. Астрофизическая модель глобального климата Земли // Киев:Наукова думка, 2003, 214 с.

[2] C. L. Badger, I. George, P. T. Grifiths, C. F. Braban, R. A. Cox, and J. P. D. Abbatt Phase transitions and hygroscopic growth of aerosol particles containing humic acid and mixtures of humic acid and ammonium sulphate // Atmos. Chem. Phys., 6, 755–768, 2006.

[3] Pidwirny, M. (2013). Atmospheric composition. Retrieved from http://www.eoearth.org/view/article/150296.

[4] А. Б. Будрецкий. Новый абсолютный минимум температуры воздуха // Информационный Бюллетень Советской Антарктической Экспедиции № 105. Ленинград, Гидрометеоиздат, 1984 г.

[5] Station Vostok // Russian Antarctic Expedition http://www.aari.aq/stations/vostok/vostok_en.html.

[6] Stepanova N. On the lowest temperatures on Earth // Monthly weather review, January, 1958, p.6-10.

[7] Stratosphere global temperature time series // NOAA, National Weather Service http://www.cpc.ncep.noaa.gov/products/stratosphere/temperature/.

[8] Devendraa Siingh, Vimlesh Pant, A K Kamra. Measurements of positive ions and air-earth current density at Maitri, Antarctica // arXiv:0905.4927 [physics.ao-ph].

[9] J. Sanders. Scientists Find Evidence of Highly Oxidizing Environment Over the South Pole // July 2001, Georgia Institute of Technology Research, http://gtresearchnews.gatech.edu/newsrelease/SPOLE.html.

[10] Good plant design and operation for onshore carbon capture installations and onshore pipelines - 2.1 General properties and uses of carbon dioxide // Energy Institute, London, September 2010 (first edition), 137p.

[11] The Dry Ice 'Snowflakes' of Mars. Analysis by Ian O'Neill // Discovery news http://news.discovery.com/space/the-foggy-carbon-dioxide-snow-of-mars-120619.html.

[12] О.В. Мороженко. Методи і результати дистанційного зондування планетних атмосфер // К. : Наук. думка, 2004. - 647 с.

[13] Dr. Chowdhury's Guide to Planet Earth (2005). "The Water Cycle". WestEd. http://www.planetguide.net/book/chapter_2/water_cycle.html

[14] Petit et. al. Vostok Ice core temperature records

[15] Pamela A. Gemery, Michael Trolier, James W. C. White. Oxygen isotope exchange between carbon dioxide and water following atmospheric sampling using glass flasks // Journal of Geophysical Research: Atmospheres Vol. 101, Issue D9, pages 14415–14420, 20 June 1996.

[16] Deniro MJ, Epstein S. Relationship between the oxygen isotope ratios of terrestrial plant cellulose, carbon dioxide, and water. Science. 1979 Apr 6;204(4388):51-3. PubMed PMID: 17816736.